

University of Groningen

## Liver magnetic resonance imaging

Sijens, Paul E.

*Published in:*  
World Journal of Gastroenterology

*DOI:*  
[10.3748/wjg.v16.i13.1558](https://doi.org/10.3748/wjg.v16.i13.1558)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2010

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Sijens, P. E. (2010). Liver magnetic resonance imaging: State of the art. *World Journal of Gastroenterology*, 16(13), 1558-1559. <https://doi.org/10.3748/wjg.v16.i13.1558>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## Liver magnetic resonance imaging: State of the art

Paul E Sijens

Paul E Sijens, Department of Radiology, University Medical Center Groningen and University of Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands

**Author contributions:** Sijens PE, series editor of the Topic Highlight “MRI of the liver, state of the art” wrote this introduction.

**Correspondence to:** Dr. Paul E Sijens, PhD, Associate Professor, Department of Radiology, University Medical Center Groningen and University of Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands. [p.e.sijens@rad.umcg.nl](mailto:p.e.sijens@rad.umcg.nl)

Telephone: +31-50-3613534 Fax: +31-50-3611798

Received: March 19, 2010 Revised: March 22, 2010

Accepted: March 29, 2010

Published online: April 7, 2010

### Abstract

Magnetic resonance imaging (MRI) has now been used for about three decades to characterize the human liver in a non-invasive way, that is without the need of using ionizing radiation or removing tissue samples. During the past few years, technical progress has been considerable and novel applications of MRI have been implemented in the clinic. The beginning of a new decade offers an excellent opportunity for having five experts to present their view on the current status of MRI (and magnetic resonance spectroscopy) in the study of perfusion, fat and iron contents, diffusion and the metabolism of diffuse liver diseases. This topic highlight series thus provides an update of current knowledge in the field of liver MRI.

© 2010 Baishideng. All rights reserved.

**Key words:** Liver; Magnetic resonance imaging

Sijens PE. Liver magnetic resonance imaging: State of the art. *World J Gastroenterol* 2010; 16(13): 1558-1559 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v16/i13/1558.htm> DOI: <http://dx.doi.org/10.3748/wjg.v16.i13.1558>

focus on five magnetic resonance imaging (MRI) methods offering opportunities for parametric exploration of the liver<sup>[1]</sup>. The first contribution focuses on the microcirculatory status in liver parenchyma and liver lesions as studied by perfusion MRI, i.e. MRI with the use of (contrast) agents to improve the contrast between the features on images<sup>[2]</sup>. Great progress has been made in turning perfusion weighted imaging into a quantitative method for detecting tumor, evaluating tumor viability after therapy and, for instance, the diagnosis of liver cirrhosis and its severity. Another important development is that nowadays, the sampling of liver tissue for the assessment of hepatic steatosis, related to alcoholic and non-alcoholic liver disease, metabolic syndrome, obesity and insuline resistance, can be replaced by the use of MRI or magnetic resonance spectroscopy (MRS) for determining liver fat content<sup>[3]</sup>. The next review is a thorough assessment of the present status in another novel application of MRI presenting an alternative to biopsy: liver iron content determination<sup>[4]</sup>. Accurate evaluation of iron overload is necessary to establish the diagnosis of hemochromatosis and guide chelation treatment in transfusion-dependent anemia. Diffusion weighted imaging (DWI), best used in combination with conventional unenhanced MRI and perfusion weighted MRI, is a promising tool used in predicting tumor responsiveness and following up on-cological treatment since DWI might be capable of detecting recurrent disease earlier than conventional imaging<sup>[5]</sup>. The final contribution features the use of MRS to study metabolism in diffuse liver diseases, diabetes and cancer<sup>[6]</sup>. Although this method still is in the preclinical stage, it is anticipated that future developments such as clinical magnets with a higher field strength (3 T, 7 T) and improved delineation of multi-component signals, will lead to intensified research in metabolic syndrome, cardiovascular disease, hepato-biliary diseases, *etc.* We believe that this issue will be of interest not only to gastroenterologists, but also to those involved in metabolic studies, cell physiology and pathology.

### REFERENCES

- 1 Sijens PE. Parametric exploration of the liver by magnetic

In this Topic Highlight “Liver MRI, state of the art”, we

- 2 **Thng CH**, Koh TS, Collins DJ, Koh DM. Perfusion magnetic resonance imaging of the liver. *World J Gastroenterol* 2010; **16**: 1598-1609
- 3 **Springer F**, Machann J, Claussen CD, Schick F, Schwenzer NF. Liver fat content determined by magnetic resonance imaging and spectroscopy. *World J Gastroenterol* 2010; **16**: 1560-1566
- 4 **Tziomalos K**, Perifanis V. Liver iron content determination by magnetic resonance imaging. *World J Gastroenterol* 2010; **16**: 1587-1597
- 5 **Kele PG**, van der Jagt EJ. Diffusion weighted imaging in the liver. *World J Gastroenterol* 2010; **16**: 1567-1576
- 6 **Dagnelie PC**, Leij-Halfwerk S. Magnetic resonance spectroscopy to study hepatic metabolism in diffuse liver diseases, diabetes and cancer. *World J Gastroenterol* 2010; **16**: 1577-1586

**S- Editor** Tian L   **L- Editor** Wang XL   **E- Editor** Ma WH